

Some other models of consensus

Vivek Borkar

October 2009

School of Technology and Computer Science
Tata Institute of Fundamental Research
Homi Bhabha Road,
Mumbai 400005, India.

Other sources of consensus-like phenomena:

1. 'Population algorithms'
2. Learning in games
3. Coordination of autonomous agents

Typically involve stochastic dynamics, analyzed by the 'o.d.e. limit'.

Example 1: Ant Colony Optimization

Intuition:

- Ants sample alternative paths to food source and deposit pheromone when they do so.
- Future ants prefer high pheromone paths.
- This builds up a reinforcement mechanism that concentrates ant population on the optimal path.

$p_i(t) :=$ probability of choosing path i at time t

$f_i(p) :=$ stationary expectation of the pheromone level on path i if $p(t) \equiv p$

Assume: $p_i > p_j, j \neq i \implies f_i(p) > f_j(p) \forall j \neq i.$

Probability increased / decreased in proportion to the excess / shortfall of average pheromone level against the population average.

Thus:

$$\dot{p}_i(t) = p_i(t)[f_i(p(t)) - \sum_j p_j(t)f_j(p_j(t))].$$

'*Replicator dynamics*' ($p_i(t) \approx$ population share of species i). Can show: If $p_i(0) > p_j(0) \forall j \neq i$, then $p_i(t) \uparrow 1$.

\implies ants concentrate on a single path.

Fact: Initial bias favors the shortest path with a probability that decreases to zero as the number of ants $\uparrow \infty$.

Example 2: Learning in repeated games:

- symmetric payoff matrix $A := [[a_{ij}]]$
- $p_i(t) :=$ the fraction of population using action i
- change from action i to j with probability

$$q_{ij}(p(t)) := \exp\left(-\beta\left(\sum_k a_{ik}p_k(t) - \sum_k a_{jk}p_k(t)\right)^+\right).$$

The '*master equation*' is

$$\dot{p}_i(t) = \sum_{j \neq i} q_{ji}(p(t)) p_j(t) - p_i(t) \sum_{j \neq i} q_{ij}(p(t))$$

$H(p) := -\frac{1}{2} \sum_{i,j} a_{ij} p_i p_j + \beta^{-1} \sum_i p_i \log(p_i)$ serves as a Liapunov function $\implies p(t) \rightarrow$ a local minimum of H .

p^* a local minimum $\iff p_i^* = Z^{-1} \exp(\beta \sum_k a_{ik} p_k^*) \forall i$.

As $\beta \uparrow \infty$, $p^* \rightarrow$ 'Nash' \hat{p} :

$$\hat{p}^T A \hat{p} \geq q^T A \hat{p} \forall q \neq \hat{p}$$

Example 3: A coordination problem:

Consider solving the scalar equation $f(x) = 0$ by the 'algorithm' $\dot{x}(t) = f(x(t))$.

Suppose $x(t) \rightarrow H := \{x : f(x) = 0\}$, assumed to be finite.

'Distributed' version: $\dot{x}_i(t) = f(x_i(t)), 1 \leq i \leq N$.

Want: $x_i(t)$ converge to the *same* element of H .

Consider

$$\dot{x}(t) = h(x(t)) := (P - I)x(t) + F(x(t)), \quad (1)$$

where $P := [[p_{ij}]]$ is an irreducible stochastic matrix and $F([x_1, \dots, x_N]^T) := [f(x_1(t)), \dots, f(x_N(t))]^T$.

Assume bounded trajectories. $\frac{\partial h_i}{\partial x_j} \geq 0$ and the positive terms form an irreducible matrix.

\implies 'cooperative system' \implies converges generically to the set $G := \{x : h(x) = 0\}$ (Hirsch).

If $x^* \in H$, $[x^*, \dots, x^*]^T \in G$.

If $x_i \in H$, $x_i \neq x_j$ for some $i \neq j$, then $x = [x_1, \dots, x_N]^T \notin G$.

It is possible that $x = [x_1, \dots, x_N]^T \in G$ with x_i 's not necessarily in H . If the system

$$\sum_j p_{ij} z_j - z_i + f(z_i) = 0 \quad \forall i,$$

has no non-constant solution, then this does not occur.

Other examples:

- genetic algorithms, particle swarm algorithms, ...
- technology adoption, herd behavior, evolution of conventions, ...
- coordination and control of robots, UAVs, mobile sensors, ...

Forces leading to consensus:

1. explicit reinforcement and / or penalty mechanism
2. 'cooperative' features such as averaging
3. symmetry of the problem.